

**Information on lecture and tutorials**

Here are a few infos on the modalities of the course "**Advanced Quantum Theory**":

- The COMPUS-ID of this course is 049160001.
- You can find detailed information on lecture and tutorials on the website of our institute:  
<https://itp3.info/aqt2324>
- You can also find detailed information on lecture and tutorials on ILIAS:  
[https://ilias3.uni-stuttgart.de/goto\\_Uni\\_Stuttgart\\_crs\\_3408956.html](https://ilias3.uni-stuttgart.de/goto_Uni_Stuttgart_crs_3408956.html)
- **Written** problems have to be handed in via ILIAS and will be corrected by the tutors. You must earn at least **50 %** of the written points to be admitted to the exam.
- **Oral** problems have to be prepared for the exercise session and will be presented by a student at the blackboard. You must earn at least **66 %** of the oral points to be admitted to the exam.
- Problems marked with an asterisk (\*) are optional and can earn you bonus points.
- If you have questions regarding the problem sets, feel free to contact your tutor at any time.

**Problem 1.1: Foundations of Quantum Mechanics**

[Written | 6 pt(s)]

ID: ex\_foundations\_of\_quantum\_mechanics:aqt2324

**Learning objective**

This problem recapitulates foundational concepts of quantum mechanics and its formalism. It is based on *Student understanding of quantum mechanics*, C. Singh, Am. J. Phys. **69**, 885 (2001); <http://dx.doi.org/10.1119/1.1365404> to prevent and/or eradicate common misconceptions. A thorough and intuitive understanding of all these concepts is essential.

We refer to a generic observable  $Q$  and its corresponding quantum mechanical operator  $\hat{Q}$ . For all of the questions, the Hamiltonian and operators  $\hat{Q}$  do not depend on time explicitly.

- The eigenvalue equation for an operator  $\hat{Q}$  is given by  $\hat{Q} |\psi_i\rangle = \lambda_i |\psi_i\rangle$ , where  $i = 1, \dots, N$ . Write an expression for  $\langle \phi | \hat{Q} | \phi \rangle$ , where  $|\phi\rangle$  is a general state, in terms of the amplitudes  $\langle \phi | \psi_i \rangle$ . 1pt(s)
- If you make measurements of a physical observable  $Q$  on a system in immediate succession, do you expect the outcome to be the same every time? Justify your answer. 1pt(s)
- If you make measurements of a physical observable  $Q$  on an ensemble of identically prepared systems which are not in an eigenstate of  $\hat{Q}$ , do you expect the outcome to be the same every time? Justify your answer. 1pt(s)
- A particle is in a one-dimensional harmonic oscillator potential. Under what conditions will the expectation value of an operator  $\hat{Q}$  depend on time if 1pt(s)

- i. the particle is initially in a momentum eigenstate?
- ii. the particle is initially in an energy eigenstate?

Justify your answer in both cases.

- e) Questions (i)-(ix) refer to the following system: An electron is in a uniform magnetic field  $B$  which is pointing in the  $z$ -direction. The Hamiltonian for the spin-degree of freedom for this system is given by  $\hat{H} = -\gamma B \hat{S}_z$ , where  $\gamma$  is the gyromagnetic ratio and  $\hat{S}_z$  is the  $z$ -component of the spin angular momentum operator. 2pt(s)

Notation:  $\hat{S}_z |\uparrow\rangle = \hbar/2 |\uparrow\rangle$ ,  $\hat{S}_z |\downarrow\rangle = -\hbar/2 |\downarrow\rangle$ .

For reference, the unnormalized eigenstates of  $\hat{S}_x$  and  $\hat{S}_y$  are given by

$$\hat{S}_x(|\uparrow\rangle \pm |\downarrow\rangle) = \pm\hbar/2(|\uparrow\rangle \pm |\downarrow\rangle) \tag{1}$$

$$\hat{S}_y(|\uparrow\rangle \pm i|\downarrow\rangle) = \pm\hbar/2(|\uparrow\rangle \pm i|\downarrow\rangle). \tag{2}$$

- i. If you measure  $S_z$  of a state  $|\chi\rangle = (|\uparrow\rangle + |\downarrow\rangle)/\sqrt{2}$ , what are the possible results, and what are their respective probabilities?
- ii. If the result of the first measurement of  $S_z$  was  $\hbar/2$ , and you immediately measure  $S_z$  again, what are the possible results, and what are their respective probabilities?
- iii. If the result of the first measurement of  $S_z$  was  $\hbar/2$ , and you immediately measure  $S_x$ , what are the possible results, and what are their respective probabilities?
- iv. What is the expectation value  $\langle \hat{S}_z \rangle$  of the state  $|\chi\rangle = (|\uparrow\rangle + |\downarrow\rangle)/\sqrt{2}$ ?
- v. If the electron is initially in an eigenstate of  $\hat{S}_x$ , does the expectation value of  $\hat{S}_y$  depend on time? Justify your answer.
- vi. If the electron is initially in an eigenstate of  $\hat{S}_x$ , does the expectation value of  $\hat{S}_z$  depend on time? Justify your answer.
- vii. If the electron is initially in an eigenstate of  $\hat{S}_z$ , does the expectation value of  $\hat{S}_x$  depend on time? Justify your answer.
- viii. If the electron is initially in an eigenstate of  $\hat{S}_z$ , does the expectation value of  $\hat{S}_z$  depend on time? Justify your answer.
- ix. If the electron is initially in an eigenstate of  $\hat{S}_x$ , does the expectation value of  $\hat{S}_x$  depend on time? Justify your answer.

**Problem 1.2: Spin  $s = 1/2$  particles**

[ Oral | 4 pt(s) ]

ID: ex\_pauli\_matrices\_aqm:aqt2324

**Learning objective**

This exercise is intended to remember some properties of Pauli matrices for spin  $s = 1/2$  particles as well as getting used to the typical algebra with Levi-Civita and Kronecker Delta symbols we routinely encounter in Quantum Mechanics.

For a particle with spin  $s = 1/2$ , the Pauli matrices can be defined as

$$\hat{\sigma}_1 = \hat{\sigma}_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \hat{\sigma}_2 = \hat{\sigma}_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \text{and} \quad \hat{\sigma}_3 = \hat{\sigma}_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (3)$$

with the following multiplication rule

$$\hat{\sigma}_i \hat{\sigma}_j = \delta_{ij} \mathbb{I}_2 + i \varepsilon_{ijk} \hat{\sigma}_k \quad \text{with} \quad i, j, k = 1, 2, 3. \quad (4)$$

The Levi-Civita symbol in a three-dimensional space is represented as

$$\varepsilon_{ijk} = \begin{cases} +1 & \text{if } (i, j, k) \text{ is } (1, 2, 3), (2, 3, 1), \text{ or } (3, 1, 2), \\ -1 & \text{if } (i, j, k) \text{ is } (3, 2, 1), (1, 3, 2), \text{ or } (2, 1, 3), \\ 0 & \text{if } i = j, \text{ or } j = k, \text{ or } k = i, \end{cases} \quad (5)$$

i.e., it takes values +1 (-1) for even (odd) permutations of the indices  $(i, j, k) = (1, 2, 3)$  or 0 if any index is repeated. The Kronecker Delta is simply defined as

$$\delta_{ij} \equiv \begin{cases} 0 & \text{for } i \neq j, \\ 1 & \text{for } i = j. \end{cases} \quad (6)$$

a) Using property (4) prove that 1pt(s)

$$(\mathbf{a} \cdot \boldsymbol{\sigma})(\mathbf{b} \cdot \boldsymbol{\sigma}) = \mathbf{a} \cdot \mathbf{b} \mathbb{I}_2 + i(\mathbf{a} \times \mathbf{b}) \cdot \boldsymbol{\sigma} \quad \text{where} \quad \mathbf{a}, \mathbf{b} \in \mathbb{R}^3. \quad (7)$$

b) Show that (i)  $[\mathbf{a} \cdot \boldsymbol{\sigma}, \mathbf{b} \cdot \boldsymbol{\sigma}] = 2i(\mathbf{a} \times \mathbf{b}) \cdot \boldsymbol{\sigma}$  and (ii)  $\text{Tr}(\mathbf{a} \cdot \boldsymbol{\sigma} \mathbf{b} \cdot \boldsymbol{\sigma}) = 2 \mathbf{a} \cdot \mathbf{b}$  with equation (7). 2pt(s)

c) Find the eigenvalues and normalized eigenfunctions of the operators  $\hat{S}_i = \hbar \hat{\sigma}_i / 2$  for  $i = x, y, z$ . 1pt(s)

**Problem 1.3: Commutators**

[Oral | 5 pt(s)]

ID: ex\_commutators\_2:aqt2324

**Learning objective**

Here we recapitulate some important commutation relations with focus on position, (linear) momentum, and angular momentum operators. As commutators are ubiquitous in quantum mechanics, manipulating them should become second nature to you.

a) Show that the following identity holds for commutators of products 1pt(s)

$$[A, BC] = B[A, C] + [A, B]C. \quad (8)$$

b) Let  $g(x)$  and  $f(p)$  be analytical functions of the position and momentum operators, respectively. Show that 1pt(s)

$$[p, g(x)] = -i\hbar \frac{d}{dx} g(x), \quad (9)$$

$$[x, f(p)] = +i\hbar \frac{d}{dp} f(p). \quad (10)$$

- c) Calculate the following commutators for canonical position and momentum operators  $x$  and  $p$  1pt(s)

$$[x, p^2], \quad [x^2, p^2], \quad [xp, p^2]. \quad (11)$$

- d) Using just the canonical commutation relations of  $x$  and  $p$ , calculate  $[L_\alpha, L_\beta]$ , where 1pt(s)  
 $L_\alpha = \varepsilon_{\alpha ij} x_i p_j$  is the  $\alpha$ -component of the angular momentum operator.

- e) Now calculate  $[L_\alpha, \mathbf{L}^2]$  and  $[L_\alpha, p^2]$ . 1pt(s)