

**Problem 4.1: Hydrogen Atom – Lowest States**

[Oral | 4 pt(s)]

ID: ex\_hydrogen\_atom\_lowest\_states:aqt2324

**Learning objective**

In this exercise, we will revisit the hydrogen atom problem and determine the size of the atoms where electrons occupy the lower states.

Consider the wave functions of the Hydrogen atom  $\psi_{n,\ell,m}$ ,

$$\psi_{n,\ell,m}(r, \theta, \varphi) = R_{n,\ell}(r)Y_{\ell m}(\theta, \varphi), \tag{1}$$

$$R_{n\ell}(r) = -N_{n\ell}(2\kappa r)^\ell e^{-\kappa r} L_{n-\ell-1}^{2\ell+1}(2\kappa r), \tag{2}$$

$$N_{n\ell} = \left(\frac{1}{a}\right)^{3/2} \frac{2}{n^2(n+\ell)!} \sqrt{\frac{(n-\ell-1)!}{(n+\ell)!}}, \tag{3}$$

where  $\kappa = 1/an$ , and  $a$  refers to the Bohr radius, and  $L_p^k$  are the Laguerre polynomials.

- a) Write explicitly the  $1s$ ,  $2s$ , and  $2p_z$  wave functions, which correspond to the set of quantum numbers  $\{n, \ell, m\} = \{1, 0, 0\}$ ,  $\{2, 0, 0\}$ , and  $\{2, 1, 0\}$ , respectively. 2<sup>pt(s)</sup>
- b) Determine the expectation value of the radial components  $r$ ,  $r^2$  and  $1/r$  in the ground state (i.e. the  $1s$  state) and the  $2p_z$  state. Deduce the size of the atom in each of these states. 2<sup>pt(s)</sup>

**Problem 4.2: One-dimensional lattice and Bloch waves**

[Oral | 8 pt(s)]

ID: ex\_1d\_bloch\_theorem:aqt2324

**Learning objective**

The purpose of this exercise is to use your knowledge of translational symmetry and the Bloch's theorem to obtain the so-called Wannier functions. This representation of Bloch states is particularly useful in the context of solid-state physics.

Consider a one-dimensional lattice with lattice sites at  $R_m$  and periodic potential  $V(r) = V(r + R_m)$  (without loss of generality the origin of the coordinate system is at either one of the lattice sites). Due to Bloch's theorem the eigenfunctions of the Hamilton operator  $H$  can be written in the form  $\psi_k(r) = e^{ikr}u_k(r)$ , where  $u_k(r)$  is lattice periodic (i.e.  $u_k(r + R_m) = u_k(r)$ ),  $k$  is located in the first Brillouin zone ( $|k| \leq \pi/a$ ) and  $a$  is the distance between two lattice sites.

- a) Show that for a lattice with  $N$  sites,  $k$  has exactly  $N$  discrete values. 2<sup>pt(s)</sup>  
**Hint:** Explore the periodic boundary conditions of  $\psi_k(r)$ , i.e.,  $\psi_k(r + L) = \psi_k(r)$  if the one-dimensional lattice has length  $L = Na$ .

- b) Using Bloch's theorem (i) show that  $E_k = E_{-k}$  and then (ii) that  $\psi_k^*(r) = \psi_{-k}(r)$ , where  $E_k$  are the corresponding energy eigenvalues of the Hamiltonian  $H$ . 2pt(s)

Now assume that the length of the lattice tends to infinity ( $L \rightarrow \infty$ ). The Bloch wave functions then obey the orthogonality relation,

$$\int dr \psi_k^*(r) \psi_{k'}(r) = \Omega_B \delta(k - k'), \tag{4}$$

where  $\Omega_B$  is the volume of the first Brillouin zone. Now, consider the expansion of the wave functions,

$$\psi_k(r) = \sum_m w(r - R_m) e^{ikR_m}, \quad \text{with } w(r - R) = \frac{1}{\Omega_B} \int_{\Omega_B} dk \psi_k(r) e^{-ikR} \tag{5}$$

where the  $R_m$ -summation runs over all lattice sites and the  $k$ -integration extends along the Brillouin zone.

- c) Without using the definition of  $w$ , show that  $\psi_k(r)$  in the given representation fulfils Bloch's theorem. 2pt(s)
- d) Show that for different  $R_m$  the functions  $w$  are orthogonal to each other (these functions, which are localized around the lattice sites  $R_m$ , are called Wannier functions). 2pt(s)

**Problem 4.3: Angular momentum and rotations**

[Written | 8 pt(s)]

ID: ex\_angular\_momentum:aqt2324

**Learning objective**

Although we already investigated the angular momentum commutation relations in the first problem list, here we will do this from a different perspective, i.e., by seeing angular momentum operators as the generators of infinitesimal rotations.

Consider the angular momentum operator  $\mathbf{L} = \mathbf{r} \times \mathbf{p}$ . In the lecture it was shown that  $\mathbf{L}$  is the infinitesimal generator of rotations such that rotations around some axis  $\mathbf{n}$  with  $\mathbf{n}^2 = 1$  about some angle  $\omega$  can be written as  $U_\omega = \exp(-i\omega\mathbf{L} \cdot \mathbf{n}/\hbar)$ . If  $U_\omega$  is the operator performing a rotation around some axis  $\omega = \omega\mathbf{n}$  in the Hilbert space, i.e.  $|\phi_\omega\rangle = U_\omega |\phi\rangle$ ; a *scalar* operator  $S$  transforms like

$$U_\omega^\dagger S U_\omega = S, \tag{6}$$

and a *vector* operator  $\mathbf{X}$  transforms like

$$U_\omega^\dagger \mathbf{X} U_\omega = R_\omega \mathbf{X}, \tag{7}$$

where  $R_\omega$  is the usual rotation matrix in three dimensions around some axis  $\omega$ .

- a) Show that for a scalar operator  $S$ ,  $[\mathbf{L}, S] = 0$ . 2pt(s)

- b) Show that for a vector operator  $\mathbf{X}$  it is  $[L_i, X_j] = i\hbar\varepsilon_{ijk}X_k$ . 2pt(s)  
**Hint:** Use the representation  $(\mathcal{R}_\omega)_{ij} = [1 - \cos(\omega)]\hat{\omega}_i\hat{\omega}_j + \cos(\omega)\delta_{ij} - \sin(\omega)\varepsilon_{ijk}\hat{\omega}_k$  for the rotation matrix and linearize (7) for small  $\omega$ .
- c) Using that  $\mathbf{r}$  and  $\mathbf{p}$  are vector operators, show that  $\mathbf{L}$  is also a vector operator. 2pt(s)  
**Hint:** Consider the components of  $U_\omega^\dagger \mathbf{r} \times \mathbf{p} U_\omega$  and show that  $U_\omega^\dagger \mathbf{r} \times \mathbf{p} U_\omega = U_\omega^\dagger \mathbf{r} U_\omega \times U_\omega^\dagger \mathbf{p} U_\omega$ .
- d) Show that  $[\mathbf{L}, \mathbf{p} \cdot \mathbf{r}] = 0$  on the one hand by explicitly calculating the commutator and on the other hand by showing that  $\mathbf{p} \cdot \mathbf{r}$  is a scalar operator. 2pt(s)