

**Problem 8.1: Density matrix**

[Oral | 5 pt(s)]

ID: ex\_density\_matrix\_2:aqt2425

**Learning objective**

The purpose of this problem is to get familiar with the concept of the density matrix (operator) and calculate some important properties.

In general, the density matrix is given by

$$\rho = \sum_i p_i |\psi_i\rangle \langle\psi_i| \tag{1}$$

where  $p_i$  is the probability to be in the state  $|\psi_i\rangle$ .

- a) Show that  $\rho^\dagger = \rho$  and  $\text{tr}(\rho) = 1$ . 1pt(s)
- b) Show that  $\text{tr}(\rho^2) \leq 1$  and equality holds for a pure state. Show also that for a pure state  $\rho^2 = \rho$ . 1pt(s)
- c) Show that the expectation value of an operator  $O$  is given by  $\langle O \rangle = \text{tr}(\rho O)$ . 1pt(s)
- d) Show that the equation of motion of the density matrix  $\rho(t)$  of a system with Hamiltonian  $H$  is given by the von Neumann equation 1pt(s)

$$i\hbar \partial_t \rho(t) = [H, \rho(t)]. \tag{2}$$

- e) Consider a canonical ensemble at temperature  $T$ . Show that for a system with Hamiltonian  $H$ , the thermal density matrix is given by 1pt(s)

$$\rho = \frac{1}{Z} \sum_n e^{-\beta E_n} |n\rangle \langle n| = \frac{1}{Z} e^{-\beta H}, \tag{3}$$

where  $\beta = \frac{1}{k_B T}$ ,  $Z = \text{tr}(e^{-\beta H})$  is the partition function, and  $H |n\rangle = E_n |n\rangle$ .

**Problem 8.2: Planck's radiation law**

[Oral | 3 pt(s)]

ID: ex\_plancks\_radiation\_law:aqt2425

**Learning objective**

In this problem, you derive Planck's radiation law of a black body which was a pioneering result in modern physics and quantum theory in particular.

First, consider a single mode of the electromagnetic field (without polarization) with Hamiltonian

$$H = \hbar\omega_k a_k^\dagger a_k. \tag{4}$$

- a) Calculate the partition sum  $Z$  and write down the thermal state  $\rho$  at temperature  $T$ . 1pt(s)
- b) Calculate the mean particle number  $\bar{n} = \langle n \rangle$  and the mean energy  $\bar{E} = \langle H \rangle$  for the thermal state  $\rho$ . 1pt(s)

In order to derive Planck's radiation law, consider a three-dimensional box of volume  $V = L^3$  with periodic boundary conditions. The Hamiltonian of the system is now given by

$$H = \sum_{\mathbf{k}, \lambda} \hbar \omega_{\mathbf{k}} a_{\mathbf{k}, \lambda}^\dagger a_{\mathbf{k}, \lambda}, \tag{5}$$

where  $\omega_{\mathbf{k}} = c|\mathbf{k}|$  and  $\lambda$  is the polarization of the mode  $\mathbf{k}$ .

- c) Based on your results in subtask b), calculate the spectral energy density  $u_\omega(T)d\omega$  and the total energy density  $u(T)$ . 1pt(s)

**Hints:**

- The system now consists of independent harmonic oscillators.
- The spectral energy density  $u_\omega d\omega$  is given by the product of the energy and the density of states in the frequency interval  $[\omega, \omega + d\omega]$ .
- In order to calculate the density of states, first calculate the mode spacing and then take the limit  $L \rightarrow \infty$ .
- $\int_0^\infty dx \frac{x^3}{e^x - 1} = \frac{\pi^4}{15}$ .

**Problem 8.3: Commutator of the electric field**

[Written | 3 pt(s)]

ID: ex\_commutator\_electric\_field:aqt2425

**Learning objective**

This problem deals with the quantized electric field. You calculate its commutator and show that it preserves causality.

In this problem, you calculate the commutator of the electric field,  $[E_i(\mathbf{r}, t), E_j(\mathbf{r}', t')]$ .

- a) In a first step, calculate the commutator  $[A_i(\mathbf{r}, t), A_j(\mathbf{r}', t')]$  of the vector potential. 1pt(s)

Start by decomposing the vector potential  $\mathbf{A}(\mathbf{r}, t)$  into its normal modes

$$\mathbf{A}(\mathbf{r}, t) = \sum_{\mathbf{k}, \lambda} \left( \frac{2\pi\hbar c^2}{V\omega_{\mathbf{k}}} \right)^{1/2} (a_{\mathbf{k}, \lambda} \boldsymbol{\epsilon}(\mathbf{k}, \lambda) e^{i(\mathbf{k}\cdot\mathbf{r} - \omega_{\mathbf{k}}t)} + \text{h.c.}), \tag{6}$$

where  $\mathbf{r}$  is the spatial coordinate,  $t$  the time,  $\mathbf{k}$  the wave vector,  $V$  the quantization volume,  $\hbar$  Planck's constant,  $c$  the speed of light, and  $\omega_{\mathbf{k}} = c|\mathbf{k}|$ .  $a_{\mathbf{k}, \lambda}$  denotes the annihilation operator for wave number  $\mathbf{k}$  and polarization  $\lambda$ ,  $\boldsymbol{\epsilon}$  is the vector of polarization.

In addition, use the completeness relation of the polarization vectors,

$$\sum_{\lambda} \epsilon_i(\mathbf{k}, \lambda) \epsilon_j^*(\mathbf{k}, \lambda) = \delta_{ij} - \frac{k_i k_j}{k^2}, \tag{7}$$

in order to bring the commutator into the following form:

$$[A_i(\mathbf{r}, t), A_j(\mathbf{r}', t')] = \partial_{ij} K(\boldsymbol{\xi}, \tau), \quad (8)$$

with  $\boldsymbol{\xi} \equiv \mathbf{r} - \mathbf{r}'$  and  $\tau \equiv t - t'$ , where  $K(\boldsymbol{\xi}, \tau)$  has to be determined.

The differential operator  $\partial_{ij}$  is defined as

$$\partial_{ij} \equiv \frac{\partial^2}{c^2} \delta_{ij} - \partial_{\xi_i} \partial_{\xi_j}. \quad (9)$$

b) Next, write the commutator for the electric field  $\mathbf{E}$  in the following form

1pt(s)

$$[E_i(\mathbf{r}, t), E_j(\mathbf{r}', t')] = -\frac{1}{c^2} \frac{\partial^2}{\partial \tau^2} [A_i(\mathbf{r}, t), A_j(\mathbf{r}', t')]. \quad (10)$$

c) Finally, transform the commutator into a form which is proportional to  $\delta(\xi^2 - c^2 \tau^2)$ .  
Discuss the physical concept behind this solution.

1pt(s)

**Hint:** It is not required to evaluate the derivatives  $\partial_{ij}$ , it is sufficient to perform the integration over  $\mathbf{k}$ . This can be done by replacing the summation  $\frac{1}{V} \sum_{\mathbf{k}} \rightarrow \int \frac{d^3 k}{(2\pi)^3}$  by an integral.