

Problem 11.1: Lifetime of “classical” atoms

[Oral | 3 pt(s)]

ID: ex_lifetime_of_classical_atoms:edyn23

Learning objective

In this problem we will calculate the lifetime of a classical atom in presence of an oscillating dipole. We will learn about the stability of the atom caused by the dipole radiation in a classical and a quantum picture.

Pretend you forgot everything you have learned in your quantum mechanics course last semester and think of the hydrogen atom as a positive central charge $+e$ with mass m_p (the proton) orbited classically by a charge with opposite sign $-e$ and mass m_e (the electron). Due to $m_p \gg m_e$ you may consider the proton stationary, $\mathbf{r}_p(t) \equiv \mathbf{0}$, and the electron's position parameterized by $\mathbf{r}_e(t) = a_0 \cos(\omega_e t) \mathbf{e}_x + a_0 \sin(\omega_e t) \mathbf{e}_y$, where a_0 is the radius of the orbit.

- Use your knowledge from classical mechanics to determine the frequency ω_e of the electron motion. 1pt(s)
- Calculate the vector potential $\mathbf{A}(\mathbf{r}, t)$ of this rotating dipole. What is the radiated power of the atom? 1pt(s)
- Estimate the lifetime of the atom. To this end assume that the radius of the orbit a_0 is given by the Bohr radius a_B . Does the result match your expectations? 1pt(s)

Problem 11.2: Rotating Quadrupole

[Oral | 3 pt(s)]

ID: ex_rotating_quadropole:edyn23

Learning objective


In this exercise we will learn about the time dependent quadrupole tensor and electromagnetic waves radiated by a rotating quadrupole. Then we will derive the angular power distribution of a rotating quadrupole and compare it with the angular power distribution of an oscillating dipole.

Consider the quadrupole setup depicted below, with two pairs of opposing charges $\pm q$ fixed at the corners of a square of size a . The square lies in the xy -plane and rotates with frequency $\omega = \omega \mathbf{e}_z$ around its center.

We will focus on the electromagnetic waves radiated by this setup.

- Write down the time-dependent charge distribution $\rho(\mathbf{x}, t)$ and calculate the quadrupole tensor 1pt(s)

$$Q_{ij}(t) = \int_{\mathbb{R}^3} d^3x \rho(\mathbf{x}, t) (3x_i x_j - |\mathbf{x}|^2 \delta_{ij}) . \quad (1)$$



For this, consider the initial condition where the $+q$ charges are located on the x -axis at time $t = 0$.

Result:

$$\begin{aligned}
 Q_{3i} &= Q_{i3} = 0, \quad i = 1, 2, 3 \\
 Q_{11} &= -Q_{22} = 3qa^2 \operatorname{Re} e^{-2i\omega t} \\
 Q_{21} &= Q_{12} = 3qa^2 \operatorname{Re} i e^{-2i\omega t}
 \end{aligned}$$

b) Show that the general expression for the angular power distribution in the far-field reads 1pt(s)

$$\frac{dP}{d\Omega} = \frac{\mu_0 c^3 k^6}{4\pi 288\pi} |\hat{\mathbf{r}} \times Q_0 \hat{\mathbf{r}}|^2 \tag{2}$$

with $\hat{\mathbf{r}} = \mathbf{r}/|\mathbf{r}|$. Here, Q_0 is the quadrupole tensor, a 3×3 amplitude matrix defined by components Q_{ij} without the oscillating factor.

Hints: The fields in the far-field approximation are given by

$$\mathbf{B} = -\frac{\mu_0}{4\pi} \frac{ik^3 c e^{ikr}}{6 r} \hat{\mathbf{r}} \times Q_0 \hat{\mathbf{r}} \quad \text{and} \quad \mathbf{E} = c \mathbf{B} \times \hat{\mathbf{r}}, \tag{3}$$

and the angular power distribution reads

$$\frac{dP}{d\Omega} = \frac{r^2}{2\mu_0} \hat{\mathbf{r}} \cdot (\mathbf{E} \times \mathbf{B}^*) . \tag{4}$$

c) Evaluate Eq. (2) with $k = 2\omega/c$ in spherical coordinates for the given setup of rotating quadrupole in the Fig. Justify why the frequency is 2ω ? Compare the result of rotating quadrupole with the angular power distribution of an oscillating *dipole*. 1pt(s)

Hint: Use the result from task a).

Problem 11.3: Spherical Bessel Functions

[Written | 5 pt(s)]

ID: ex_spherical_bessel_functions:edyn23

Learning objective

The spherical Bessel and Hankel functions j_l and $h_l^{(1)} \equiv h_l^+$ play a crucial role for the expansion of the vector potential. In this exercise we will derive explicit expressions for these functions.

The spherical Bessel equation describes the radial part $R_l(r)$ of the solution $\Phi(r, \theta, \varphi) = R_l(r)Y_{lm}(\theta, \varphi)$ of the Helmholtz equation $[\Delta + k^2]\Phi = 0$ and reads

$$\left[\partial_x^2 + \frac{2}{x} \partial_x + \left(1 - \frac{l(l+1)}{x^2} \right) \right] R_l(x) = 0 \quad \text{for } l \in \mathbb{N}_0 \quad (5)$$

with $x = kr$.

- a) As a warm-up, show that for half integer $\nu = l + \frac{1}{2}$ the substitution $R_l(x) = \frac{u_l(x)}{\sqrt{x}}$ converts the spherical Bessel equation to the ordinary Bessel equation 1pt(s)

$$\left[\partial_x^2 + \frac{1}{x} \partial_x + \left(1 - \frac{\nu^2}{x^2} \right) \right] u_l(x) = 0. \quad (6)$$

Provide solutions of Eq. (5) in terms of the Bessel and Neumann functions $J_\nu(x)$ and $N_\nu(x)$ which have been introduced in the lecture during the discussion of electrostatics.

The solutions derived from $J_\nu(x)$ and $N_\nu(x)$ are denoted as $j_l(x)$ and $n_l(x)$ and referred to as *spherical* Bessel and Neumann functions, respectively (up to normalizing factors). In the remainder of this exercise, you will derive explicit expressions for these functions.

- b) To this end, prove that the *spherical Hankel functions* 1pt(s)

$$h_l^\pm(x) = \mp \frac{(x/2)^l}{l!} \int_{\pm 1}^{i\infty} dt e^{ixt} (1-t^2)^l \quad (7)$$

are solutions of Eq. (5) for $x > 0$ and $l \in \mathbb{N}_0$.

Hints: Use $x^{-1} \partial_x^2 x = 2x^{-1} \partial_x + \partial_x^2$ and write the integrand as a total derivative with respect to t .

- c) Now show that h_l^\pm satisfy the recursion relation 1pt(s)

$$\frac{dh_l^\pm(x)}{dx} = \frac{l}{x} h_l^\pm(x) - h_{l+1}^\pm(x). \quad (8)$$

- d) The spherical Hankel functions are a basis of the two-dimensional solution space for every l . 1pt(s)
Another common basis is given by the linear combinations

$$j_l(x) = \frac{1}{2} [h_l^+(x) + h_l^-(x)] \quad \text{and} \quad n_l(x) = \frac{1}{2i} [h_l^+(x) - h_l^-(x)] \quad (9)$$

which are the *spherical* Bessel and Neumann functions as introduced in task a).

Use the recursion from task c) to prove the explicit expressions

$$j_l(x) = (-x)^l \left(\frac{1}{x} \frac{d}{dx} \right)^l \frac{\sin(x)}{x} \quad (10a)$$

$$n_l(x) = -(-x)^l \left(\frac{1}{x} \frac{d}{dx} \right)^l \frac{\cos(x)}{x} \quad (10b)$$

These are known as *Rayleigh's formulas*.

Hint: Use mathematical induction.

- e) Use the above results to write down $j_l(x)$, $n_l(x)$ and $h_l^+(x)$, $h_l^-(x)$ for $l = 0, 1$ and sketch the graphs of $j_l(x)$, $n_l(x)$. 1^{pt(s)}