

**Problem 2.1: Green's function for the Poisson equation**

[ Oral | 3 pt(s) ]

ID: ex\_greens\_function\_poisson\_equation:edyn23

**Learning objective**

In this problem, we prove the important identity of Green function in two ways. First with Gauss law and second by taking Fourier transformation of Yukawa potential in the zero mass limit.

Consider the following important identity

$$\Delta \frac{1}{|\mathbf{r}|} = -4\pi \delta^3(\mathbf{r}) \quad (1)$$

- a) Show that  $\Delta \frac{1}{|\mathbf{r}|} = 0$  for any  $\mathbf{r} \neq 0$ . Then, use Gauss's law to prove (1) by integrating over a sphere. 1pt(s)

The Yukawa potential is given by the expression

$$\phi_Y(\mathbf{r}) = \frac{e^{-mr}}{r} \quad (2)$$

where  $r = |\mathbf{r}|$  and  $m > 0$  is the mass of the particle that mediates the potential. The inverse mass is proportional to a length scale (Compton length) that determines the range of the potential. If photons had a rest mass, the Coulomb potential would have to be replaced by the Yukawa potential. We can see that the Coulomb potential is the limiting case of  $\phi_Y(\mathbf{r})$  in the zero-mass limit (infinite-range limit).

- b) Use the three-dimensional Fourier transformation of  $\Delta \frac{1}{|\mathbf{r}|}$  to show that equation (1) holds. First calculate the Fourier transform of  $\frac{1}{|\mathbf{r}|}$  by transforming the Yukawa potential and then take the zero-mass limit. 1pt(s)
- c) Calculate the Fourier transformation of the equation 1pt(s)

$$[\Delta - m^2] \phi_Y(\mathbf{r}) = -4\pi \delta^3(\mathbf{r}), \quad (3)$$

and evaluate the solution in Fourier space for  $\phi_Y(\mathbf{k})$ . Determine  $\phi_Y(\mathbf{r})$  by explicitly evaluating the Fourier transformation and demonstrate that the solution is the Yukawa potential. **Hint:** Use the result from (b) to obtain the solution in Fourier space. The Fourier transformation into real space can be achieved using the residue theorem.

**Problem 2.2: Cavendish experiment, part 1: Spherical capacitor**

[Oral | 3 pt(s)]

ID: ex\_cavendish\_experiment\_part1:edyn23

**Learning objective**

In this problem, we will learn to calculate the electric field, scalar potential and capacitance of a spherical capacitor which is made of two concentric metallic spherical shells.

A spherical capacitor is given by two concentric, metallic spherical shells with radii  $R_1 < R_2$  and respective charges  $Q_1 = Q$  and  $Q_2 = -Q$ .

- Calculate the electric field  $\mathbf{E}(\mathbf{r})$  of the given arrangement for the three regions  $r < R_1$ ,  $R_1 < r < R_2$  and  $R_2 < r$ . 1pt(s)
- Determine the scalar potential  $\phi(\mathbf{r})$  with the boundary condition  $\phi(\mathbf{r}) \rightarrow 0$  for  $r \rightarrow \infty$  such that it is continuous at  $R_1$  and  $R_2$ . 1pt(s)
- Determine the capacitance of the spherical capacitor. 1pt(s)

**Problem 2.3: Cavendish experiment, part 2**

[Written | 2 (+1 bonus) pt(s)]

ID: ex\_cavendish\_experiment\_part2:edyn23

**Learning objective**

In this problem, we will calculate the total energy of the spherical capacitor by calculating the electrostatic interaction energy. We also calculate the charge distribution on both spheres.

In 1772, Cavendish designed the following experiment to verify Coulombs law. He used a spherical capacitor with the two spheres initially connected by an electrical contact. He then placed a static charge on the outer sphere and subsequently removed the contact between the spheres. After removing the outer sphere, he confirmed that the inner sphere had not been charged. This is a special property of the  $1/r$  Coulomb law where the electric potential inside a conducting sphere is constant (the electric field vanishes).

- Assume that the electrostatic potential is instead given by the modified power law  $\phi_\epsilon(\mathbf{r}) = \frac{1}{|\mathbf{r}|^{1-\epsilon}}$ . Determine the charge on the inner sphere for both prospective potentials as follows. Let  $V_i$  be the “volume” of the  $i$ th sphere. We can calculate the electrostatic interaction energy by 1pt(s)

$$E_{ij} = \frac{1}{2} \int_{V_i} d^3r \int_{V_j} d^3r' \rho(\mathbf{r}) \rho(\mathbf{r}') \phi(\mathbf{r} - \mathbf{r}'), \quad (4)$$

where  $E_{ii}$  is the self energy of the  $i$ th sphere and  $E_{12} + E_{21} = 2E_{12}$  is the interaction energy between the two spheres. The full energy is given by  $E = E_{11} + E_{22} + 2E_{12}$ .

- Due to the spherical symmetry, the charge density will be homogeneously distributed on the spheres. Assume that the spheres have a surface density of  $\sigma_i = Q_i/S_i$  were  $S_i$  is the surface area of the respective sphere. Calculate the total energy  $E(Q_1, Q_2)$  for arbitrary charges.

- ii) Determine the actual charge distribution by minimizing the energy with the constraint that the total charge  $Q = Q_1 + Q_2$  is fixed.
- b) Repeat the analysis assuming that electrostatic potential is given by the Yukawa potential with a finite photon mass. 1pt(s)
- \*c) How can we derive the capacitance from Problem 2.2c from the energy function  $E(Q_1, Q_2)$  for the Coulomb potential (set  $\epsilon = 0$  or  $m = 0$  in your solution). +1pt(s)