

Prof. Dr. Hans-Peter Büchler
 Institute for Theoretical Physics III, University of Stuttgart

May 8th, 2024
 SS 2024

Problem 5.1: Potential of an infinitely long cylinder

[Oral | 2 pt(s)]

ID: ex_potential_infinite_cylinder:edyn24

Learning objective

Here, we solve Laplace's equation in cylindrical coordinates with Dirichlet boundary conditions.

We consider an infinitely long, hollow cylinder of radius R . Using Laplace's equation in cylindrical coordinates, we determine the electric potential inside and outside of the cylinder, given the value of the potential on the boundary of the cylinder.

- a) Assume the potential on the boundary is given by

1pt(s)

$$\phi(z, \varrho = R, \varphi) = \phi_0 + \phi_1 \cos \varphi, \quad (1)$$

where z is the axial coordinate, φ is the polar angle, and ϱ the radial distance in cylindrical coordinates. Think about the geometry of the problem and calculate the potential inside and outside of the cylinder.

- b) The potential on the boundary of the cylinder is

1pt(s)

$$\phi(z, \varrho = R, \varphi) = \cos(kz) (\phi_0 + \phi_1 \cos \varphi), \quad (2)$$

with $k \neq 0$. Calculate the potential and determine its value in the limit $\varrho \rightarrow \infty$ (for taking this limit, it is helpful to look up the asymptotic behavior of the Bessel function e.g. on Wikipedia).

Problem 5.2: Electric field of a dipole

[Written | 6 pt(s)]

ID: ex_electric_field_dipole:edyn24

Learning objective

In the first part of the problem, we calculate the electric field for a dipole. The resulting expression contains a δ -function term, whose physical importance is discussed in the second part of the problem.

- a) Recall the important result $\Delta \frac{1}{|\mathbf{r}|} = -4\pi \delta^3(\mathbf{r})$ from Problem 2.1 and generalize it to

1pt(s)

$$\partial_\alpha \partial_\beta \frac{1}{|\mathbf{r}|} = -\frac{\delta_{\alpha\beta}}{|\mathbf{r}|^3} + 3\frac{x_\alpha x_\beta}{|\mathbf{r}|^5} - \frac{4\pi}{3} \delta_{\alpha\beta} \delta^3(\mathbf{r}). \quad (3)$$

Hint: Use a symmetry argument and the result from exercise Problem 2.1 to derive the last term in equation (3).

- b) In the lecture, it was demonstrated that the electric potential for a dipole \mathbf{p} is given by $\phi(\mathbf{r}) = \frac{\mathbf{p} \cdot \mathbf{r}}{4\pi\epsilon_0 |\mathbf{r}|^3} = -(\mathbf{p} \cdot \nabla) \frac{1}{4\pi\epsilon_0 |\mathbf{r}|}$. Using relation (3), show that the electric field of the dipole can be written as ($\hat{\mathbf{r}} = \mathbf{r}/|\mathbf{r}|$): 1pt(s)

$$\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \left[\frac{3(\hat{\mathbf{r}} \cdot \mathbf{p}) \hat{\mathbf{r}} - \mathbf{p}}{|\mathbf{r}|^3} - \frac{4\pi}{3} \mathbf{p} \delta^3(\mathbf{r}) \right]. \quad (4)$$

The δ -function term in equation (4) is a correction for $\mathbf{r} = 0$. In the following, we are going to re-derive it in a different way to understand its physical origin.

We would like to prove the following THEOREM: The *average electric field* over the volume V enclosed by a sphere of radius R , due to an arbitrary charge distribution within the sphere, is given by

$$\overline{\mathbf{E}} = -\frac{1}{4\pi\epsilon_0} \frac{\mathbf{p}}{R^3}, \quad (5)$$

where \mathbf{p} is the total dipole moment with respect to the center of the sphere.

- c) To do this, first calculate the average electric field within the sphere (with enclosed volume V), due to a single charge q at position \mathbf{r}_q : 1pt(s)

$$\overline{\mathbf{E}}_q = \frac{1}{V} \int_V d^3r \mathbf{E}_q(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \frac{q}{V} \int_V d^3r \frac{\mathbf{r} - \mathbf{r}_q}{|\mathbf{r} - \mathbf{r}_q|^3}. \quad (6)$$

Realize that this expression can also be considered as the electric field *at the position* \mathbf{r}_q , that is generated by a (fictional) ball with a uniform charge density $\rho = -q/V$. Use this analogy to calculate $\overline{\mathbf{E}}_q$ via Gauss's law.

- d) Use the superposition principle to generalize the result for the point charge q to arbitrary charge distributions and prove equation (5). 1pt(s)
- e) Explicitly calculate the average electric field that is generated by a point-like dipole, by integrating the electric field from equation (4) over a ball. In your integration, start by excluding a small region around the origin. 1pt(s)
- f) Finally, show that the δ -function term in equation (5) is essential to satisfy the average-value theorem. 1pt(s)

Note: Another approach is to calculate the electric field of a homogeneously polarized ball of radius a . Outside of the ball, the field is exactly given by equation (4). Inside the ball, the field has a constant value $\mathbf{E}_{\text{in}} = -1/4\pi\epsilon_0 \cdot \mathbf{p}/a^3$, where \mathbf{p} is the dipole moment of the ball. As the size of the ball goes to zero, the field strength goes to infinity in such a way that the integral over the ball remains constant, giving the prefactor of the δ -function: $-\mathbf{p}/3\epsilon_0$.

Problem 5.3: Spherical multipole moment


[Oral | 2 pt(s)]

ID: ex_spherical_multipole_moment:edyn24

Learning objective

The goal of this problem is to calculate the spherical multipole moments q_{lm} for different charge distributions and to study when a quadrupole moment occurs.

We perform calculations for two charge distributions (A) and (B). Both consist of four charges in the xy -plane, placed distance a from the origin and equidistant to each other. The distributions are given in the sketch



- a) Write down the charge distribution in spherical coordinates. The relation between the charge distribution in Cartesian coordinates $\rho(x, y, z)$ and spherical coordinates $\rho_{\text{sph}}(r, \theta, \phi)$ is given by (why?): 1pt(s)

$$\rho(x, y, z) = \frac{\rho_{\text{sph}}(r, \theta, \phi)}{r^2 \sin \theta} \quad (7)$$

- b) Compute the spherical monopole, dipole and quadrupole moments for both arrangements. 1pt(s)