

Problem 9.1: Gaussian disorder

[Oral | 4 (+2 bonus) pt(s)]

ID: ex_gaussian_impurity:qft24

Learning objective

In this exercise, we will once again go through the replica trick introduced in the lecture and apply it to the calculation of the quasiparticle's lifetime in the presence of the Gaussian disorder.

Let us start by splitting the action into two parts,

$$S = S_c[\psi^\dagger, \psi] + S_d[\psi^\dagger, \psi, V],$$

where the S_c describes the clean electron system, and S_d corresponds to the coupling between electrons and disorder potential,

$$S_d[\psi^\dagger, \psi, V] = \sum_{\omega_n} \int d^d r \psi^\dagger(\omega_n, \mathbf{r}) V(\mathbf{r}) \psi(\omega_n, \mathbf{r}).$$

We will consider the disorder potential to be Gaussian, namely,

$$\langle V(\mathbf{r}) V(\mathbf{r}') \rangle_V = \frac{\gamma^2}{(2\pi s^2)^{d/2}} e^{-|\mathbf{r}-\mathbf{r}'|^2/(2s^2)}, \quad (1)$$

here we define the averaging over the disorder $\langle \cdot \rangle_V$ as

$$\langle \mathcal{A}[V] \rangle_V = \mathcal{N} \int D[V] \mathcal{A}[V] \exp \left[-\frac{1}{L^{2d}} \frac{(2\pi s^2)^{d/2}}{2\gamma^2} \int d^d r \int d^d r' V(\mathbf{r}) e^{|\mathbf{r}-\mathbf{r}'|^2/(2s^2)} V(\mathbf{r}') \right],$$

where \mathcal{N} is the normalization constant. Then the average value of a fermionic operator \mathcal{O} reads

$$\begin{aligned} \langle \langle \mathcal{O} \rangle_S \rangle_V &= \int D[V] P[V] \int D[\psi^\dagger, \psi] \mathcal{O}[\psi^\dagger, \psi] e^{-S_c[\psi^\dagger, \psi] - S_d[\psi^\dagger, \psi, V]}, \\ P[V] &= \mathcal{N} \exp \left[-\frac{1}{L^{2d}} \frac{(2\pi s^2)^{d/2}}{2\gamma^2} \int d^d r \int d^d r' V(\mathbf{r}) e^{|\mathbf{r}-\mathbf{r}'|^2/(2s^2)} V(\mathbf{r}') \right] \end{aligned}$$

Even though we have already set up everything to use the perturbation theory straightforwardly, let us proceed with the *replica trick*. The basic idea behind the replica trick is to introduce R identical copies of the considered system, average over all the copies and finally take the limit $R \rightarrow 0$,

$$\langle \langle \mathcal{O} \rangle_S \rangle_V = \lim_{R \rightarrow 0} \frac{1}{R} \int D[V] P[V] \int D[\psi^\dagger, \psi] \sum_{a=1}^R \mathcal{O}[\psi_a^\dagger, \psi_a] e^{-\sum_{a=1}^R S[\psi_a^\dagger, \psi_a, V]}. \quad (2)$$

Suppose we want to find the current induced by the external homogeneous electric field $\mathbf{F}(\mathbf{r}, t) = \mathbf{E}$. First, linearize the kinetic equation with respect to the electric field, assuming that $f_{\mathbf{p}}(\mathbf{r}, t) \approx f_0(\varepsilon_{\mathbf{p}}) + \delta f_{\mathbf{p}}$ with $f_0(\varepsilon_{\mathbf{p}}) = (1 + e^{\beta(\varepsilon_{\mathbf{p}} - \mu)})^{-1}$, $\delta f_{\mathbf{p}} \sim E$, and $I_{\text{col}}[f_0] = 0$, $I_{\text{col}}[\delta f_{\mathbf{p}}] = -\delta f_{\mathbf{p}}/\tau_{\mathbf{p}}$, where $\tau_{\mathbf{p}}$ is the life-time found earlier. *Try to come up with an intuitive explanation of why the time change of the distribution function related to the collision with impurities should be related to the lifetime.* Then calculate the current density to obtain the Drude formula.